

KÄHLER MANIFOLDS ADMITTING A FLAT COMPLEX CONFORMAL CONNECTION

G. GANCHEV AND V. MIHOVA

ABSTRACT. We prove that any Kähler manifold admitting a flat complex conformal connection is a Bochner-Kähler manifold with special scalar distribution and zero geometric constants. Applying the local structural theorem for such manifolds we obtain a complete description of the Kähler manifolds under consideration.

1. INTRODUCTION

Let (M, g, J) ($\dim M = 2n$) be a Kähler manifold with complex structure J , metric g , Levi-Civita connection ∇ , curvature tensor R , Ricci tensor ρ and scalar curvature τ . The Bochner curvature tensor $B(R)$ is given by

$$\begin{aligned} B(R)(X, Y)Z &= R(X, Y)Z \\ &- Q(Y, Z)X + Q(X, Z)Y - g(Y, Z)Q(X) + g(X, Z)Q(Y) \\ &- Q(JY, Z)JX + Q(JX, Z)JY + 2Q(JX, Y)JZ \\ &- g(JY, Z)JQ(X) + g(JX, Z)JQ(Y) + 2g(JX, Y)JQ(Z), \quad X, Y, Z \in \mathfrak{X}M, \end{aligned}$$

where $Q(X, Y) = \frac{1}{2(n+2)}\rho(X, Y) - \frac{\tau}{8(n+1)(n+2)}g(X, Y)$ and $Q(X)$ is the corresponding tensor of type (1.1).

The manifold is said to be *Bochner flat* (*Bochner-Kähler*) if its Bochner curvature tensor vanishes identically, i.e.

$$\begin{aligned} (1.1) \quad R(X, Y)Z &= \\ &Q(Y, Z)X - Q(X, Z)Y + g(Y, Z)Q(X) - g(X, Z)Q(Y) \\ &+ Q(JY, Z)JX - Q(JX, Z)JY - 2Q(JX, Y)JZ \\ &+ g(JY, Z)JQ(X) - g(JX, Z)JQ(Y) - 2g(JX, Y)JQ(Z), \quad X, Y, Z \in \mathfrak{X}M, \end{aligned}$$

For any real C^∞ function u on M we denote $\omega = du$ and $P = \text{grad } u$.

In [10] Yano introduced on a Kähler manifold a complex conformal connection and proved

Theorem A. *If in a $2n$ -dimensional ($n \geq 2$) Kähler manifold there exists a scalar function u such that the complex conformal connection*

$$\begin{aligned} \mathcal{D}_X Y &= \nabla_X Y + \omega(X)Y + \omega(Y)X - g(X, Y)P \\ &- \omega(JX)JY - \omega(JY)JX - g(JX, Y)JP, \quad X, Y \in \mathfrak{X}M, \end{aligned}$$

is of zero curvature, then the Bochner curvature tensor of the manifold vanishes.

1991 *Mathematics Subject Classification.* Primary 53B35, Secondary 53C25.

Key words and phrases. Complex conformal connection, Bochner-Kähler manifolds with special scalar distribution, warped product Kähler manifolds.

In [7] Seino proved the inverse

Theorem B. *In a Kählerian space with vanishing Bochner curvature tensor if there exists a non-constant function u satisfying the equality*

$$(\nabla_X \omega)(Y) + 2\omega(JX)\omega(JY) + \omega(P)g(X, Y) = 0,$$

then the complex conformal connection is of zero curvature.

In this paper we prove

Theorem 3.1. *A Kähler manifold (M, g, J) ($\dim M = 2n \geq 6$) admits a flat complex conformal connection if and only if it is a Bochner-Kähler manifold whose scalar distribution D_τ is a B_0 -distribution with function $a + k^2 = 0$ and geometric constants $\mathfrak{B} = \mathfrak{b}_0 = 0$.*

Applying the local structural theorem [2] for Bochner-Kähler manifolds whose scalar distribution is a B_0 -distribution, we describe locally all Kähler manifolds admitting a flat complex conformal connection.

2. PRELIMINARIES

Let (M, g, J) ($\dim M = 2n$) be a Kähler manifold with metric g , complex structure J and Levi-Civita connection ∇ . We denote by $\mathfrak{X}M$ the Lie algebra of all $\mathcal{C}^\infty$ vector fields on M . The fundamental Kähler form Ω is defined as follows

$$\Omega(X, Y) = g(JX, Y), \quad X, Y \in \mathfrak{X}M.$$

For any $\mathcal{C}^\infty$ real function u on M we consider the conformal metric $\bar{g} = e^{2u}g$. We denote the 1-form $\omega := du$ and $P := \text{grad } u$ with respect to the metric g . Then $(M, \bar{g}, J)$ is a locally conformal Kähler manifold, or a W_4 -manifold in the classification scheme of [3]. The fundamental Kähler form and the Lee form of the structure $(\bar{g}, J)$ are $\bar{\Omega}(X, Y) = \bar{g}(JX, Y)$, $X, Y \in \mathfrak{X}M$ and $\bar{\omega} = 2\omega = 2du$, respectively. The Lee vector $\bar{P}$ corresponding to $\bar{\omega}$ with respect to the metric $\bar{g}$ is $\bar{P} = 2e^{-2u}P$.

The unique linear connection $\mathcal{D}$ with torsion $\mathcal{T}$ satisfying the conditions:

$$\begin{aligned} (2.1) \quad & 1) \mathcal{D}J = 0; \\ & 2) \mathcal{D}\bar{g} = 0; \\ & 3) \mathcal{T} = -\bar{\Omega} \otimes J\bar{P} \end{aligned}$$

is said to be a *complex conformal connection* [10].

In terms of the Kähler structure (g, J) $\mathcal{D}$ is given by

$$\begin{aligned} (2.2) \quad \mathcal{D}_X Y = & \nabla_X Y + \omega(X)Y + \omega(Y)X - g(X, Y)P \\ & - \omega(JX)JY - \omega(JY)JX - g(JX, Y)JP, \quad X, Y \in \mathfrak{X}M. \end{aligned}$$

The conditions (2.1) in terms of the Kähler structure (g, J) become

$$\begin{aligned} (2.3) \quad & 1) \mathcal{D}J = 0; \\ & 2) \mathcal{D}g = -2\omega \otimes g; \\ & 3) \mathcal{T} = -2\Omega \otimes JP. \end{aligned}$$

Denote by $\mathcal{R}$ the curvature tensor of the complex conformal connection $\mathcal{D}$. Taking into account (2.2) we have the relation between R and $\mathcal{R}$:

$$\begin{aligned}
(2.4) \quad & \mathcal{R}(X, Y)Z = R(X, Y)Z \\
& -\{(\nabla_Y \omega)(Z) - \omega(Y)\omega(Z) + \omega(JY)\omega(JZ) + \frac{1}{2}\omega(P)g(Y, Z)\}X \\
& +\{(\nabla_X \omega)(Z) - \omega(X)\omega(Z) + \omega(JX)\omega(JZ) + \frac{1}{2}\omega(P)g(X, Z)\}Y \\
& -g(Y, Z)\{\nabla_X P - \omega(X)P - \omega(JX)JP + \frac{1}{2}\omega(P)X\} \\
& +g(X, Z)\{\nabla_Y P - \omega(Y)P - \omega(JY)JP + \frac{1}{2}\omega(P)Y\} \\
& +\{(\nabla_Y \omega)(JZ) - \omega(Y)\omega(JZ) - \omega(JY)\omega(Z) + \frac{1}{2}\omega(P)g(Y, JZ)\}JX \\
& -\{(\nabla_X \omega)(JZ) - \omega(X)\omega(JZ) - \omega(JX)\omega(Z) + \frac{1}{2}\omega(P)g(X, JZ)\}JY \\
& +g(Y, JZ)\{\nabla_X JP - \omega(X)JP + \omega(JX)P + \frac{1}{2}\omega(P)JX\} \\
& -g(X, JZ)\{\nabla_Y JP - \omega(Y)JP + \omega(JY)P + \frac{1}{2}\omega(P)JY\} \\
& -(\nabla_X \omega)(JY)JZ + (\nabla_Y \omega)(JX)JZ + 2g(X, JY)\{\omega(JZ)P + \omega(Z)JP\}
\end{aligned}$$

for all $X, Y, Z \in \mathfrak{X}M$.

From (2.4) it follows that the curvature tensor $\mathcal{R}$ satisfies the first Bianchi identity (i.e. $\mathcal{R}$ is a Kähler tensor) if and only if [7]:

$$(2.5) \quad (\nabla_X \omega)(Y) + 2\omega(JX)\omega(JY) + \omega(P)g(X, Y) = 0, \quad X, Y \in \mathfrak{X}M,$$

which is equivalent to the condition

$$\mathcal{D}_X P = 0, \quad X \in \mathfrak{X}M.$$

If the 1-form ω satisfies (2.5), then (2.4) becomes

$$\begin{aligned}
(2.6) \quad & \mathcal{R}(X, Y)Z = R(X, Y)Z \\
& +L(Y, Z)X - L(X, Z)Y + g(Y, Z)L(X) - g(X, Z)L(Y) \\
& +L(JY, Z)JX - L(JX, Z)JY - 2L(JX, Y)JZ \\
& +g(JY, Z)JL(X) - g(JX, Z)JL(Y) - 2g(JX, Y)JL(Z), \quad X, Y, Z \in \mathfrak{X}M,
\end{aligned}$$

where $L(X, Y) = \omega(X)\omega(Y) + \omega(JX)\omega(JY) + \frac{1}{2}\omega(P)g(X, Y)$ and $L(X)$ is the corresponding tensor of type (1,1) with respect to the Kähler metric g .

If (M, g, J) admits a flat complex conformal connection (2.2), then $\mathcal{R}$ satisfies the first Bianchi identity, i.e. (2.5) holds good. Then (2.6) implies that the Kähler manifold is Bochner flat.

Conversely, if (M, g, J) admits a 1-form ω satisfying (2.5), then (2.4) becomes (2.6). The condition (M, g, J) is Bochner flat implies that $\mathcal{R} = 0$, i.e. the complex conformal connection (2.2) is flat.

3. A CURVATURE CHARACTERIZATION OF KÄHLER MANIFOLDS ADMITTING FLAT COMPLEX CONFORMAL CONNECTION

For any Bochner-Kähler manifold (M, g, J) in [2] we proved that

$$(3.1) \quad (\nabla_X \rho)(Y, Z) = \frac{1}{4(n+1)} \{2d\tau(X)g(Y, Z) + d\tau(Y)g(X, Z) + d\tau(Z)g(X, Y) + d\tau(JY)g(X, JZ) + d\tau(JZ)g(X, JY)\}, \quad X, Y, Z \in \mathfrak{X}M.$$

This equality shows that the conditions $\tau = \text{const}$ and $\nabla \rho = 0$ are equivalent on a Bochner-Kähler manifold. Because of the structural theorem in [8] the case $B(R) = 0$, $d\tau = 0$, can be considered as well-studied.

We consider Bochner-Kähler manifolds satisfying the condition $d\tau \neq 0$ for all points $p \in M$. This condition allows us to introduce the frame field

$$\left\{ \xi = \frac{\text{grad } \tau}{\|d\tau\|}, \quad J\xi = \frac{J \text{grad } \tau}{\|d\tau\|} \right\}$$

and the J -invariant distributions D_τ and $D_\tau^\perp = \text{span}\{\xi, J\xi\}$.

Thus our approach to the local theory of Bochner-Kähler manifolds is to treat them as Kähler manifolds (M, g, J, D_τ) endowed with a J -invariant distribution D_τ generated by the Kähler structure (g, J) . We call this distribution *the scalar distribution* of the manifold [2].

A J -invariant distribution D_τ , $(D_\tau^\perp = \text{span}\{\xi, J\xi\})$ is said to be a B_0 -distribution [1] if $\dim M = 2n \geq 6$ and

$$\begin{aligned} i) \quad & \nabla_{x_0} \xi = \frac{k}{2} x_0, \quad k \neq 0, \quad x_0 \in D_\tau, \\ ii) \quad & \nabla_{J\xi} \xi = -p^* J\xi, \\ iii) \quad & \nabla_\xi \xi = 0, \end{aligned}$$

where k and p^* are functions on M .

The above conditions are equivalent to the equalities

$$(3.2) \quad \begin{aligned} \nabla_X \xi &= \frac{k}{2} \{X - \eta(X)\xi + \eta(JX)J\xi\} + p^* \eta(JX)J\xi, \quad X \in \mathfrak{X}M, \\ dk &= \xi(k)\eta, \quad p^* = -\frac{\xi(k) + k^2}{k}. \end{aligned}$$

In [2] we have shown that

$$(3.3) \quad \mathfrak{B} = \|\rho\|^2 - \frac{\tau^2}{2(n+1)} + \frac{\Delta\tau}{n+1}$$

is a constant on any Bochner-Kähler manifold. We call this constant *the Bochner constant* of the manifold.

Let us denote

$$\begin{aligned}
4\pi(X, Y)Z &:= g(Y, Z)X - g(X, Z)Y - 2g(JX, Y)JZ \\
&\quad + g(JY, Z)JX - g(JX, Z)JY, \\
8\Phi(X, Y)Z &:= g(Y, Z)(\eta(X)\xi - \eta(JX)J\xi) - g(X, Z)(\eta(Y)\xi - \eta(JY)J\xi) \\
&\quad + g(JY, Z)(\eta(X)J\xi + \eta(JX)\xi) - g(JX, Z)(\eta(Y)J\xi + \eta(JY)\xi) \\
&\quad - 2g(JX, Y)(\eta(Z)J\xi + \eta(JZ)\xi) \\
&\quad + (\eta(Y)\eta(Z) + \eta(JY)\eta(JZ))X - (\eta(X)\eta(Z) + \eta(JX)\eta(JZ))Y \\
&\quad - (\eta(Y)\eta(JZ) - \eta(JY)\eta(Z))JX + (\eta(X)\eta(JZ) - \eta(JX)\eta(Z))JY \\
&\quad + 2(\eta(X)\eta(JY) - \eta(JX)\eta(Y))JZ, \quad X, Y, Z \in \mathfrak{X}M.
\end{aligned}$$

In [2] we have also proved that

If (M, g, J) ($\dim M = 2n \geq 6$) is a Bochner-Kähler manifold whose scalar distribution D_τ is a B_0 -distribution, then

$$(3.4) \quad R = a\pi + b\Phi, \quad b \neq 0,$$

where a, b are the following functions on M

$$(3.5) \quad a = \frac{\tau}{(n+1)(n+2)} + \frac{2b_0}{n+2}, \quad b = \frac{2\tau}{(n+1)(n+2)} - \frac{2nb_0}{n+2},$$

and

$$(3.6) \quad b_0 = \frac{2a-b}{2} = \text{const.}$$

In [2] we studied three classes of Bochner-Kähler manifolds whose scalar distribution is a B_0 -distribution according to the function $a + k^2$:

$$a + k^2 > 0, \quad a + k^2 = 0, \quad a + k^2 < 0.$$

Now we can prove a curvature characterization of Kähler manifolds admitting a flat complex conformal connection.

Theorem 3.1. *A Kähler manifold (M, g, J) ($\dim M = 2n \geq 6$) admits a flat complex conformal connection if and only if it is a Bochner-Kähler manifold whose scalar distribution D_τ is a B_0 -distribution with function $a + k^2 = 0$ and geometric constants $\mathfrak{B} = \mathfrak{b}_0 = 0$.*

Proof. Let u be a $\mathcal{C}^\infty$ function on M , such that the complex conformal connection $\mathcal{D}$, given by (2.2) with $\omega = du \neq 0$, $P = \text{grad } u$, is flat. Then (2.5) and (2.6) imply that the curvature tensor R of M has the structure (1.1). Comparing the tensor Q from (1.1) and the tensor L from (2.6) we obtain

$$(3.7) \quad \rho(X, Y) = -2(n+2)\{\omega(X)\omega(Y) + \omega(JX)\omega(JY) + \omega(P)g(X, Y)\}, \quad X, Y \in \mathfrak{X}M$$

and

$$(3.8) \quad \rho(X, P) = -2(n+2)\omega(P)\omega(X), \quad X \in \mathfrak{X}M.$$

After taking a trace in (3.7) we also get

$$(3.9) \quad \tau = -4(n+1)(n+2)\omega(P).$$

Taking into account (2.5) we calculate from (3.7)

$$(3.10) \quad (\nabla_X \rho)(Y, Z) = 2(n+2) \omega(P) \{2\omega(X)g(Y, Z) + \omega(Y)g(X, Z) + \omega(Z)g(X, Y) \\ + \omega(JY)g(X, JZ) + \omega(JZ)g(X, JY)\}, \quad X, Y, Z \in \mathfrak{X}M.$$

Comparing (3.1) and (3.10) in view of (3.9), we obtain

$$(3.11) \quad \omega = -\frac{d\tau}{2\tau}, \quad P = -\frac{\text{grad } \tau}{2\tau}, \quad \|d\tau\|^2 = \frac{-\tau^3}{(n+1)(n+2)}.$$

The unit vector field $\xi = \frac{\text{grad } \tau}{\|d\tau\|}$ because of (3.11) gets the form

$$\xi = 2\sqrt{\frac{(n+1)(n+2)}{-\tau}} P.$$

From (2.5) and (3.9) we obtain

$$(3.12) \quad \nabla_X \xi = -\frac{1}{2} \sqrt{\frac{-\tau}{(n+1)(n+2)}} \{X - \eta(X)\xi - 2\eta(JX)J\xi\}, \quad X \in \mathfrak{X}M.$$

Now from (3.2) and (3.12) it follows that the scalar distribution D_τ of the manifold is a B_0 -distribution with functions

$$(3.13) \quad k = -\sqrt{\frac{-\tau}{(n+1)(n+2)}}, \quad p^* = \frac{3}{2} \sqrt{\frac{-\tau}{(n+1)(n+2)}}.$$

Then (2.6) and (3.11) give that the curvature tensor R of the manifold has the form

$$R = \frac{\tau}{(n+1)(n+2)} (\pi + 2\Phi)$$

and the functions a and b are

$$(3.14) \quad a = \frac{\tau}{(n+1)(n+2)}, \quad b = \frac{2\tau}{(n+1)(n+2)}.$$

From (3.13) and (3.14) we find $a + k^2 = 0$. The equalities (3.6) and (3.14) imply that $\mathfrak{b}_0 = 0$.

Taking into account (2.5), (3.11) and (3.7) we find

$$(3.15) \quad \Delta \tau = \frac{-\tau^2}{n+1}, \quad \|\rho\|^2 = \frac{(n+3)\tau^2}{2(n+1)^2}.$$

Replacing $\Delta \tau$ and $\|\rho\|^2$ in (3.3) we obtain $\mathfrak{B} = 0$.

For the inverse, let (M, g, J) be a Bochner-Kähler manifold whose scalar distribution is a B_0 -distribution. Then it follows [2] that (3.2), (3.4) and (3.5) hold good. Under the condition $\mathfrak{b}_0 = 0$ we find that the functions a and b satisfy (3.14).

The condition $a + k^2 = 0$ implies that $k^2 = -a = \frac{-\tau}{(n+1)(n+2)}$, i.e. $\tau < 0$. From Theorem 3.5 in [1] it follows that

$$\|d\tau\| = \xi(\tau) = \frac{(n+1)(n+2)}{2} \xi(b) = \frac{(n+1)(n+2)}{2} kb > 0,$$

which gives that the function k is negative and

$$k = -\sqrt{\frac{-\tau}{(n+1)(n+2)}}, \quad \|d\tau\|^2 = \frac{-\tau^3}{(n+1)(n+2)}, \quad p^* = \frac{3}{2} \sqrt{\frac{-\tau}{(n+1)(n+2)}}.$$

Then, from the equality (3.2) for any $X, Y \in \mathfrak{X}M$ we have

$$(\nabla_X \eta)(Y) = -\frac{1}{2} \sqrt{\frac{-\tau}{(n+1)(n+2)}} \{g(X, Y) - \eta(X)\eta(Y) + 2\eta(JX)\eta(JY)\}.$$

Putting $2u := -\ln(-\tau)$ and $\omega := du = -\frac{d\tau}{2\tau} = -\frac{\|d\tau\|}{2\tau} \eta = -\frac{k}{2} \eta$ we prove that $(\nabla_X \omega)(Y)$ satisfies (2.5) and the complex conformal connection (2.2) is flat. **QED**

Let $(Q_0, g_0, \varphi, \tilde{\xi}_0, \tilde{\eta}_0)$ be an α_0 -Sasakian space form [4] with constant φ -holomorphic sectional curvatures H_0 . In [2] we introduced warped product Kähler manifolds, which are completely determined by the underlying α_0 -Sasakian space form Q_0 of type $H_0 + 3\alpha_0^2 \begin{smallmatrix} \geq \\ \leq \end{smallmatrix} 0$ and the generating function $p(t)$, $t \in I \subset \mathbb{R}$.

In order to obtain a local description of the Kähler manifolds admitting a flat complex conformal connection we apply Theorem 6.1 in [2], which states:

Any Bochner-Kähler manifold whose scalar distribution is a B_0 -distribution locally has the structure of a warped product Kähler manifold with generating function $p(t)$ (or $t(p)$) of type 1. – 13.

According to Theorem 3.1 any Kähler manifold (M, g, J) , ($\dim M = 2n \geq 6$) admitting a flat complex conformal connection is locally a Bochner-Kähler manifold whose scalar distribution is a B_0 -distribution with function $a + k^2 = 0$ and constants $\mathfrak{B} = \mathfrak{b}_0 = 0$. In terms of [2] the conditions $\mathfrak{B} = \mathfrak{b}_0 = 0$ are equivalent to the conditions $\mathfrak{K} = \mathfrak{b}_0 = 0$.

Hence, (M, g, J) is a warped product Bochner-Kähler manifold whose underlying α_0 -Sasakian space form is of type $H_0 + 3\alpha_0^2 = 0$ with metric

$$g = p^2(t) \left\{ g_0 + \left(\frac{1}{\alpha_0} \frac{dp}{dt} - 1 \right) \tilde{\eta}_0 \otimes \tilde{\eta}_0 \right\} + \eta \otimes \eta,$$

generated by the function

$$p(t) = \frac{1}{\sqrt[3]{1 - 3\alpha_0(t - t_0)}}, \quad t \in \left(-\infty, \frac{1 + 3\alpha_0 t_0}{3\alpha_0} \right)$$

of type 9. [2].

This metric is not complete.

Especially in the case $\alpha_0 = 1$ the underlying manifold is a Sasakian space form with $H_0 = -3$. Sasakian space forms of type $H_0 = -3$ have been studied by Ogiue [5] and Okumura [6]. A classification theorem for Sasakian space forms under the assumption of completeness has been given by Tanno [9].

REFERENCES

- [1] G. Ganchev, V. Mihova, Kähler manifolds of quasi-constant holomorphic sectional curvatures, *ArXiv: math.DG/0505671*, to appear.
- [2] G. Ganchev, V. Mihova, Warped product Kähler manifolds and Bochner-Kähler metrics, *ArXiv: math.DG/0605082*, to appear.
- [3] A. Gray, L. Hervella, The sixteen classes of almost Hermitian manifolds and their linear invariants, *Ann. di Mat. Pura ed Appl.*, **123**(1980), 35-58.
- [4] D. Janssens, L. Vanhecke, Almost contact structures and curvature tensors, *Kodai Math. J.*, **4** (1981), 1-27.
- [5] K. Ogiue, On almost contact manifolds admitting axiom of planes or axiom of free mobility, *Kodai Math. Sem. Rep.*, **16**(1964), 223-232.

- [6] M. Okumura, On infinitesimal conformal and projective transformations of normal contact spaces, *Tôhoku Math. J.*, **14**(1962), 398-412.
- [7] M. Seino, Some considerations on various curvature tensors, *Hokkaido Math. J.*, **10**(1981), 13-26.
- [8] S. Tachibana, R. C. Liu, Notes on Kählerian metrics with vanishing Bochner curvature tensor, *Kodai Math. Sem. Rep.*, **22** (1970), 313-321.
- [9] S. Tanno, Sasakian manifolds with constant φ -holomorphic sectional curvature, *Tôhoku Math. J.*, **21**(1969), 501-507.
- [10] K. Yano, On complex conformal connections, *Kodai Math. Sem. Rep.*, **26**(1975), 137-151.

BULGARIAN ACADEMY OF SCIENCES, INSTITUTE OF MATHEMATICS AND INFORMATICS, ACAD. G. BONCHEV STR. BL. 8, 1113 SOFIA, BULGARIA
E-mail address: ganchev@math.bas.bg

FACULTY OF MATHEMATICS AND INFORMATICS, UNIVERSITY OF SOFIA, J. BOUCHIER STR. 5, (1164) SOFIA, BULGARIA
E-mail address: mihova@fmi.uni-sofia.bg